Author Index

Abbiati, G., see Fossati, T. 577(1992)382
Abe, H., see O'Dowd, A. 577(1992)347
Addison, T. E., see Hopkins, N. K. 577(1992)87
Altieri, I., see Pichini, S. 577(1992)358
Alvarez, E., see Saunders, P. P. 577(1992)37
Alvarez, J. G. and Touchstone, J. C.
Separation of acidic and neutral lipids by aminopropyl-bonded silica gel column chromatography 577(1992)142
Andersen, J. V. and Hansen, S. H.
Simultaneous quantitative determination of naproxen, its metabolite 6-O-desmethylnaproxen and their five conjugates in plasma and urine samples by high-performance liquid chromatography on dynamically modified silica 577(1992)325
Andersen, J. V. and Hansen, S. H.
Simultaneous determination of (*R*)- and (*S*)-naproxen and (*R*)- and (*S*)-6-O-desmethylnaproxen by high-performance liquid chromatography on a Chiral-AGP column 577(1992)362
Andersen, M. P., see Jørgensen, M. 577(1992)167
Anderson, C. B., see Mangino, M. J. 577(1992)157
Baird, W. M., see Teffera, Y. 577(1992)69
Barbhaiya, R. H., see Janiszewski, J. S. 577(1992)151
Barré, J., see Hervé, F. 577(1992)43
Bavazzano, P., see Busini, S. 577(1992)180
Beetens, J., see Weber, C. 577(1992)1
Bennett, S. M., see Jajoo, H. K. 577(1992)299
Berdecia-Rodriguez, J., see Hendrich, C. E. 577(1992)19
Berger, M. R., see Gebelein, M. 577(1992)146
Blair, I. A., see Edeki, T. 577(1992)190
Bouquillon, A. I., Freeman, D. and Moulin, D. E.
Simultaneous solid-phase extraction and chromatographic analysis of morphine and hydromorphone in plasma by high-performance liquid chromatography with electrochemical detection 577(1992)354
Brisson, J., see Hopkins, N. K. 577(1992)87
Brocks, D. R., see Liang, W. T. C. 577(1992)317
Busini, S., Bavazzano, P. and Occhiato, E.
High-performance liquid chromatographic determination of 3,5-dimethylhippuric acid in the occupational exposure to trimethylbenzenes 577(1992)180
Campestrini, J., see Rouan, M. C. 577(1992)387
Castiglioni, C., see Fossati, T. 577(1992)382
Chollet, D. and Künstner, P.
Fast systematic approach for the determination of drugs in biological fluids by fully automated high-performance liquid chromatography with on-line solid-phase extraction and automated cartridge exchange. Application to cebaracetam in human urine 577(1992)335
Christensen, J. V.
Determination of (+)-5-(2,3-dihydrobenzofuran-7-yl)-3-methyl-8-nitro-2,3,4,5-tetrahydro-1H-3-benzazepin-7-ol (NNC 01-0687), a novel dopamine D-1 receptor antagonist, in plasma by solid-phase extraction and high-performance liquid chromatography 577(1992)366
Dandekar, K. A., see Janiszewski, J. S. 577(1992)151
De Clerck, F., see Weber, C. 577(1992)1
Delhotal Landes, B., Miscoria, G. and Flouvat, B.
Determination of lansoprazole and its metabolites in plasma by high-performance liquid chromatography using a loop column 577(1992)117
Duché, J.-C., see Hervé, F. 577(1992)43
Edeki, T., Robin, D. W., Prakash, C., Blair, I. A. and Wood, A. J. J.
Sensitive assay for triazolam in plasma following low oral doses 577(1992)190
Escarzaga, M., see Watts, M. T. 577(1992)289
Favier, A., see Richard, M. J. 577(1992)9
Ferris, R., Missbichler, A., Hartter, E., Mann-Buxbaum, E., Schalkhammer, T., Urban, G. and Pittner, F.
Construction and use of two α-human atrial natriuretic peptide-fragment affinity chromatography columns in the isolation of C- and N-terminal epitope-specific antibodies for use in a prototype α-hANP biosensor 577(1992)251
Filippich, L. J., see Zhu, J. 577(1992)77
Flouvat, B., see Delhotal Landes, B. 577(1992)117
Fossati, T., Parisi, S., Abbiati, G. and Castiglioni, C.
Determination of picotamide in human plasma and urine by high-performance liquid chromatography 577(1992)382
Frantsuzova, N. A., see Klyashchitsky, B. A. 577(1992)267
Freeman, D., see Bouquillon, A. I. 577(1992)354
Friedrich, A., see Hagedorn, H.-W. 577(1992)195
Gaillard, Y., Gay-Montchamp, J. P. and Ollagnier, M.
Gas chromatographic determination of meprobamate in serum or plasma after solid-phase extraction 577(1992)171
Gaspari, F., Taiocchi, L. and Pochobradsky, M. G.
Determination of brodimoprim and its hydroxy metabolite in human plasma, blood and urine by high-performance liquid chromatography 577(1992)123
Gay-Montchamp, J. P., see Gaillard, Y. 577(1992)171
Gebelein, M., Merdes, G. and Berger, M. R.
Nucleotide preparation from cells and determination of nucleotides by ion-pair high-performance liquid chromatography 577(1992)146
Geodakyan, S. V., Voskoboinikova, I. V., Kolesnik, J. A., Tjukavkina, N. A., Litvinenko, V. I. and Glyzin, V. I.
High-performance liquid chromatographic method for the determination of mangiferin, likviritin and dihydroquercetin in rat plasma and urine 577(1992)371
Glyzin, V. I., see Geodakyan, S. V. 577(1992)371
Godbillon, J., see Rouan, M. C. 577(1992)387
Guiraud, P., see Richard, M. J. 577(1992)9
Hage, D. S., see Noctor, T. A. G. 577(1992)305
Hagedorn, H.-W., Schulz, R. and Friedrich, A.
Detection of methandienone (methandrostenolone) and metabolites in horse urine by gas chromatography–mass spectrometry 577(1992)195
Haginaka, J. and Seyama, C.
Determination of enantiomers of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride (E2020), a centrally acting acetylcholine esterase inhibitor, in plasma by liquid chromatography with fluorometric detection 577(1992)95

Hansen, S. H., see Andersen, J. V. 577(1992)325
Hansen, S. H., see Andersen, J. V. 577(1992)362
Hansen, S. H., see Jørgensen, M. 577(1992)167
Hartter, E., see Ferris, R. 577(1992)251
Hendrich, C. E., Berdecia-Rodriguez, J., Wiedmeier, V. T. and Porterfield, S. P.
Method for the quantitation of iodothyronines in body tissues and fluids using high-performance liquid chromatography 577(1992)19
Heo, G. S., see Lee, K.-J. 577(1992)135
Hervé, F., Duché, J.-C., Barré, J., Millot, M.-C. and Tillement, J.-P.
pH titration curves of the desialylated human α_1-acid glycoprotein variants by combined isoelectrofocusing–electrophoresis: utilization in the development of a fractionation method for the protein variants by chromatography on immobilized metal affinity adsorbent 577(1992)43
Hirabayashi, N., see Ikegawa, S. 577(1992)229
Hirata, H., see Okutani, R. 577(1992)25
Hopkins, N. K., Wagner, C. M., Brisson, J. and Addison, T. E.
Validation of the simultaneous determination of methylprednisolone and methylprednisolone acetate in human plasma by high-performance liquid chromatography 577(1992)87
Hung, M.-F., see Kastrissios, H. 577(1992)103
Ikegawa, S., Hirabayashi, N., Yoshimura, T., Tohma, M., Maeda, M. and Tsuji, A.
Determination of conjugated bile acids in human urine by high-performance liquid chromatography with chemiluminescence detection 577(1992)229
Itoh, Y., see Okutani, R. 577(1992)25
Jajoo, H. K., Bennett, S. M. and Kornhauser, D. M.
Thermospray liquid chromatographic–mass spectrometric analysis of anti-AIDS nucleosides: quantification of 2′,3′-dideoxycytidine in plasma samples 577(1992)299
Jamali, F., see Liang, W. T. C. 577(1992)317
Janiszewski, J. S., Mulvana, D. E., Kaul, S., Dandekar, K. A. and Barbhaiya, R. H.
High-performance liquid chromatographic determination of 2′,3′-didehydro-3′-deoxythymidine, a new anti-human immunodeficiency virus agent, in human plasma and urine 577(1992)151
Jørgensen, M., Andersen, M. P. and Hansen, S. H.
Simultaneous determination of nitroglycerin and its dinitrate metabolites by capillary gas chromatography with electron-capture detection 577(1992)167
Kantarjian, H. M., see Saunders, P. P. 577(1992)37
Kapil, R. P., Padovani, P. K., King, S.-Y. P. and Lam, G. N.
Nanogram level quantitation of oxycodone in human plasma by capillary gas chromatography using nitrogen–phosphorus selective detection 577(1992)283
Kasahara, T., see Okutani, R. 577(1992)25
Kastrissios, H., Hung, M.-F. and Triggs, E. J.
High-performance liquid chromatographic method for the quantitation of bupivacaine, 2,6-pipecoloxylidide and 4′-hydroxybµpivacaine in plasma and urine 577(1992)103
Kaul, S., see Janiszewski, J. S. 577(1992)151
Kawai, T., see Okutani, R. 577(1992)25
Kelly, M. T., see Sagar, K. A. 577(1992)109
Kim, N. J., see Lee, K.-J. 577(1992)135
King, S.-Y. P., see Kapil, R. P. 577(1992)283
Klyashchitsky, B. A., Mitina, V. K., Frantsuzova, N. A., Krasnopolsky, Y. M. and Shvets, V. I.
Use of biospecific interactions of collagen, fibronectin and their fragments in affinity chromatography 577(1992)267
Kolesnik, J. A., see Geodakyan, S. V. 577(1992)371
Kornhauser, D. M., see Jajoo, H. K. 577(1992)299
Krasnopolsky, Y. M., see Klyashchitsky, B. A. 577(1992)267
Kumps, A., see Mardens, Y. 577(1992)341
Künstner, P., see Chollet, D. 577(1992)335
Lam, G. N., see Kapil, R. P. 577(1992)283
Lecaillon, J. B., see Rouan, M. C. 577(1992)387
Lee, K.-J., Heo, G. S., Kim, N. J. and Moon, D.-C.
Separation of theophylline and its analogues by micellar electrokinetic chromatography: application to the determination of theophylline in human plasma 577(1992)135
Liang, W. T. C., Brocks, D. R. and Jamali, F.
Stereospecific high-performance liquid chromatographic assay of pirprofen enantiomers in rat plasma and urine 577(1992)317
Litvinenko, V. I., see Geodakyan, S. V. 577(1992)371
Liu, H. and Wehmeyer, K. R.
Solid-phase extraction with supercritical fluid elution as a sample preparation technique for the ultratrace analysis of flavone in blood plasma 577(1992)61
Lowes, S., Rose, M. E., Mills, G. A. and Pollitt, R. J.
Identification of urinary acylcarnitines using gas chromatography–mass spectrometry: preliminary clinical applications 577(1992)205
MacFarlane, N., see O'Dowd, A. 577(1992)347
Maeda, M., see Ikegawa, S. 577(1992)229
Mangino, M. J., Zografakis, J., Murphy, M. K. and Anderson, C. B.
Improved and simplified tissue extraction method for quantitating long-chain acyl-coenzyme A thioesters with picomolar detection using high-performance liquid chromatography 577(1992)157
Mann-Buxbaum, E., see Ferris, R. 577(1992)251
Mardens, Y., Kumps, A., Planchon, C. and Wurth, C.
Comparison of two extraction procedures for urinary organic acids prior to gas chromatography–mass spectrometry 577(1992)341
Matuszewski, B., see Woolf, E. 577(1992)129
Maylin, G., see Steffenrud, S. 577(1992)221
Meo, J., see Richard, M. J. 577(1992)9
Merdes, G., see Gebelein, M. 577(1992)146
Miller, D. J., see O'Dowd, A. 577(1992)347
Millot, M.-C., see Hervé, F. 577(1992)43
Mills, G. A., see Lowes, S. 577(1992)205
Miscoria, G., see Delhotal Landes, B. 577(1992)117
Missbichler, A., see Ferris, R. 577(1992)251
Mitina, V. K., see Klyashchitsky, B. A. 577(1992)267
Moncrieff, J.
Extractionless determination of diclofenac sodium in serum using reversed-phase high-performance liquid chromatography with fluorimetric detection 577(1992)185

Monger, L. S. and Olliff, C. J.
Measurement of β-endorphin in human plasma by high-performance liquid chromatography with electrochemical detection: validation of a method employing the simultaneous purification and concentration of β-endorphin 577(1992)239
Moon, D.-C., see Lee, K.-J. 577(1992)135
Mori, H., see Nagasawa, M. 577(1992)275
Motohashi, M., see Yamashita, K. 577(1992)174
Moulin, D. E., see Bouquillon, A. I. 577(1992)354
Mukaida, N., see Okutani, R. 577(1992)25
Mulvana, D. E., see Janiszewski, J. S. 577(1992)151
Murphy, M. K., see Mangino, M. J. 577(1992)157
Nagasawa, M., Sasabe, H., Shimizu, T. and Mori, H.
Use of a carrier for quantitation of a new dihydropyridine calcium antagonist (OPC-13340) in human plasma by highly sensitive gas chromatography with negative-ion chemical ionization mass spectrometry 577(1992)275
Ng, J., see Zhu, J. 577(1992)77
Noctor, T. A. G., Wainer, I. W. and Hage, D. S.
Allosteric and competitive displacement of drugs from human serum albumin by octanoic acid, as revealed by high-performance liquid affinity chromatography, on a human serum albumin-based stationary phase 577(1992)305
Occhiato, E., see Busini, S. 577(1992)180
O'Dowd, A., O'Dowd, J. J., O'Dowd, J. J. M., MacFarlane, N., Abe, H. and Miller, D. J.
Analysis of novel imidazoles from isolated perfused rabbit heart by two high-performance liquid chromatographic methods 577(1992)347
O'Dowd, J. J., see O'Dowd, A. 577(1992)347
O'Dowd, J. J. M., see O'Dowd, A. 577(1992)347
Ohmori, S., see Otsuka, M. 577(1992)215
Okutani, R., Itoh, Y., Hirata, H., Kasahara, T., Mukaida, N. and Kawai, T.
Simple and high-yield purification of urine protein 1 using immunoaffinity chromatography: evidence for the identity of urine protein 1 and human Clara cell 10-kilodalton protein 577(1992)25
Ollagnier, M., see Gaillard, Y. 577(1992)171
Olliff, C. J., see Monger, L. S. 577(1992)239
Otsuka, M. and Ohmori, S.
Simple and sensitive determination of diacetyl and acetoin in biological samples and alcoholic drinks by gas chromatography with electron-capture detection 577(1992)215
Ottaviani, G., see Pichini, S. 577(1992)358
Pacifici, R., see Pichini, S. 577(1992)358
Padovani, P. K., see Kapil, R. P. 577(1992)283
Parisi, S., see Fossati, T. 577(1992)382
Parviainen, M. T., see Pikkarainen, S. A. 577(1992)163
Pichini, S., Altieri, I., Pacifici, R., Rosa, M., Ottaviani, G. and Zuccaro, P.
Simultaneous determination of cotinine and *trans*-3′-hydroxycotinine in human serum by high-performance liquid chromatography 577(1992)358
Pikkarainen, S. A. and Parviainen, M. T.
Determination of retinyl palmitate and total vitamin A content in liver and liver-based ready-to-eat foods 577(1992)163
Pittner, F., see Ferris, R. 577(1992)251
Planchon, C., see Mardens, Y. 577(1992)341
Pochobradsky, M. G., see Gaspari, F. 577(1992)123
Pollitt, R. J., see Lowes, S. 577(1992)205
Porterfield, S. P., see Hendrich, C. E. 577(1992)19
Prakash, C., see Edeki, T. 577(1992)190
Richard, M. J., Guiraud, P., Meo, J. and Favier, A.
High-performance liquid chromatographic separation of malondialdehyde–thiobarbituric acid adduct in biological materials (plasma and human cells) using a commercially available reagent 577(1992)9
Robin, D. W., see Edeki, T. 577(1992)190
Rosa, M., see Pichini, S. 577(1992)358
Rose, M. E., see Lowes, S. 577(1992)205
Rouan, M. C., Campestrini, J., Lecaillon, J. B. and Godbillon, J.
Rapid determination of propyphenazone in plasma by high-performance liquid chromatography 577(1992)387
Sagar, K. A., Kelly, M. T. and Smyth, M. R.
Analysis of terbutaline in human plasma by high-performance liquid chromatography with electrochemical detection using a micro-electrochemical flow cell 577(1992)109
Sasabe, H., see Nagasawa, M. 577(1992)275
Saunders, P. P., Alvarez, E. and Kantarjian, H. M.
Determination of nicotinamide-adenine dinucleotide and thiazole-4-carboxamide-adenine dinucleotide in human leukocytes by reversed-phase high-performance liquid chromatography 577(1992)37
Schalkhammer, T., see Ferris, R. 577(1992)251
Schulz, R., see Hagedorn, H.-W. 577(1992)195
Seyama, C., see Haginaka, J. 577(1992)95
Shearer, G., see Tarbin, J. A. 577(1992)376
Shimizu, T., see Nagasawa, M. 577(1992)275
Shvets, V. I., see Klyashchitsky, B. A. 577(1992)267
Smith, D. L., see Teffera, Y. 577(1992)69
Smyth, M. R., see Sagar, K. A. 577(1992)109
Steffenrud, S. and Maylin, G.
Thermospray liquid chromatography–mass spectrometry of corticosteroids 577(1992)221
Taiocchi, L., see Gaspari, F. 577(1992)123
Tarbin, J. A. and Shearer, G.
High-performance liquid chromatographic determination of imidocarb in cattle kidney with cation-exchange clean-up 577(1992)376
Teffera, Y., Baird, W. M. and Smith, D. L.
Detection of benzo[*a*]pyrene sulfate and glucuronide conjugates in cell culture medium by directly coupled microbore high-performance liquid chromatography–fast atom bombardment mass spectrometry 577(1992)69
Tegtmeier, F., see Weber, C. 577(1992)1
Tillement, J.-P., see Hervé, F. 577(1992)43
Tjukavkina, N. A., see Geodakyan, S. V. 577(1992)371
Tohma, M., see Ikegawa, S. 577(1992)229
Touchstone, J. C., see Alvarez, J. G. 577(1992)142
Triggs, E. J., see Kastrissios, H. 577(1992)103
Tsuji, A., see Ikegawa, S. 577(1992)229
Urban, G., see Ferris, R. 577(1992)251
Voskoboinikova, I. V., see Geodakyan, S. V. 577(1992)371
Wagner, C. M., see Hopkins, N. K. 577(1992)87
Wainer, I. W., see Noctor, T. A. G. 577(1992)305

Watts, M. T., Escarzaga, M. and Williams, C. H.
Gas chromatographic headspace analysis of sevoflurane in blood 577(1992)289
Weber, C., Beetens, J., De Clerck, F. and Tegtmeier, F.
Gas chromatographic–mass spectrometric determination of 11-dehydrothromboxane B_2 in human urine 577(1992)1
Wehmeyer, K. R., see Liu, H. 577(1992)61
Wiedmeier, V. T., see Hendrich, C. E. 577(1992)19
Williams, C. H., see Watts, M. T. 577(1992)289
Wood, A. J. J., see Edeki, T. 577(1992)190
Woolf, E. and Matuszewski, B.
High-performance liquid chromatographic procedure for the determination of a non-nucleoside HIV-1 reverse transcriptase inhibitor in human plasma 577(1992)129
Wurth, C., see Mardens, Y. 577(1992)341
Yamashita, K., Motohashi, M. and Yashiki, T.
Automated high-performance liquid chromatographic method for the simultaneous determination of cefotiam and Δ^3-cefotiam in human plasma using column switching 577(1992)174
Yashiki, T., see Yamashita, K. 577(1992)174
Yoshimura, T., see Ikegawa, S. 577(1992)229
Zhu, J., Ng, J. and Filippich, L. J.
Determination of tannic acid and its phenolic metabolites in biological fluids by high-performance liquid chromatography 577(1992)77
Zografakis, J., see Mangino, M. J. 577(1992)157
Zuccaro, P., see Pichini, S. 577(1992)358